

Natural Statistics and Wildlife Probability

Statistics & Probability · Practice Test · 25 Questions

1. In a population of honeybees, the foraging range follows a distribution where the mean distance from the hive is approximately 2 to 3 kilometers. What statistical term describes the most frequently occurring value in this distance dataset?

- A) Mean
- B) Median
- C) Mode
- D) Standard Deviation

2. If a biologist tags 50 sea turtles in a bay and later recaptures 200, finding 10 with tags, what is the estimated total population size using the Lincoln-Peterson index formula?

- A) 500
- B) 1,000
- C) 2,000
- D) 2,500

3. The gestation period of African elephants has a normal distribution. If the mean is 660 days with a standard deviation of 15 days, what percentage of births fall within one standard deviation of the mean?

- A) 68%
- B) 95%
- C) 99.7%
- D) 50%

4. When calculating the probability of a bird species carrying a specific genetic mutation in a population where 1 in 50 individuals is affected, what is the probability that a randomly selected individual does NOT carry the mutation?

- A) 0.02
- B) 0.98
- C) 0.05
- D) 0.95

5. In a forest ecosystem, the frequency of tree heights often follows a specific statistical curve. Which distribution is most commonly used to model the number of occurrences of rare events, such as the sighting of a specific nocturnal predator in an hour?

- A) Normal
- B) Poisson
- C) Binomial
- D) Exponential

6. If the probability of a sea lion pup surviving its first year is 0.75, what is the probability of two independent pups both surviving their first year?

- A) 0.5625
- B) 1.50
- C) 0.25
- D) 0.80

7. Which statistical measure represents the square root of the variance in the annual rainfall data of the Amazon rainforest?

- A) Range
- B) Mean
- C) Standard Deviation
- D) Interquartile Range

8. In a study of 100 emperor penguins, 60 were male and 40 were female. What is the probability of selecting a female penguin at random?

- A) 0.6
- B) 0.4
- C) 0.5
- D) 0.2

9. The lifespan of a certain species of wildflower follows an exponential distribution. In this type of probability distribution, what is the relationship between the mean and the standard deviation?

- A) Mean equals standard deviation
- B) Mean is double the standard deviation
- C) Mean is half the standard deviation
- D) They are unrelated

10. If a coral reef survey shows that the number of fish species present increases logarithmically with the area sampled, which statistical relationship is being observed?

- A) Linear regression
- B) Species-area relationship
- C) Normal distribution
- D) Chi-square test

11. In a sample of 200 butterflies, the wingspans are measured. If the mean wingspan is 50mm and the median is 50mm, what does this suggest about the distribution of the data?

- A) It is heavily skewed
- B) It is perfectly symmetrical
- C) It is bimodal
- D) It has high outliers

12. What is the probability of picking an egg that will hatch into a female alligator if the sex ratio of the clutch is 1:1?

- A) 0.25
- B) 0.5
- C) 0.75
- D) 1.0

13. When researchers study the biodiversity of a soil sample, they use the Shannon Diversity Index. What does a higher value in this index statistically represent?

- A) Lower species count
- B) Higher species richness and evenness
- C) Decreased environmental stability
- D) A smaller sample size

14. If a population of wolves has a growth rate that is constant over time, which type of function best represents the total population size over years?

- A) Linear
- B) Exponential
- C) Logarithmic
- D) Inverse

15. In a data set of 1,000 birds, 250 are migratory. What is the relative frequency of migratory birds in this population?

- A) 0.025
- B) 0.25
- C) 2.5
- D) 25

16. Which statistical test is commonly used to compare the observed frequencies of animal diet choices against expected frequencies based on prey availability?

- A) T-test
- B) Chi-square test
- C) ANOVA
- D) Correlation coefficient

17. The body mass of a species of lizard is found to have a coefficient of variation of 10%. If the mean mass is 200g, what is the standard deviation?

- A) 10g
- B) 20g
- C) 30g
- D) 40g

18. When tracking the migration path of a whale, the distance covered per day fluctuates. What statistical measure would best indicate the 'typical' daily distance while minimizing the effect of extreme outliers?

- A) Mean
- B) Median
- C) Range
- D) Sum

19. In a box plot showing the distribution of nesting depths for turtles, what does the box itself represent?

- A) The range
- B) The interquartile range
- C) The mean
- D) The standard deviation

20. If there is a 5% chance that a rare orchid will bloom in a specific year, what is the probability it will NOT bloom?

- A) 0.05
- B) 0.95
- C) 0.5
- D) 0.99

21. Which term describes the data point that lies furthest from the mean in a dataset of animal migration speeds?

- A) Mode
- B) Outlier
- C) Median
- D) Quartile

22. In a population of 500 insects, 50 are predators. What is the probability of randomly choosing a non-predatory insect?

- A) 0.1
- B) 0.9
- C) 0.5
- D) 0.09

23. What is the sum of all probabilities in a sample space representing the possible weather conditions for a habitat (Sunny, Rainy, Cloudy)?

- A) 0
- B) 0.5
- C) 1
- D) 100

24. A researcher measures the concentration of pollutants in a stream. What statistical value is calculated by dividing the sum of all concentrations by the number of samples?

- A) Median
- B) Mode
- C) Mean
- D) Variance

25. If the correlation coefficient between local temperature and butterfly emergence date is -0.85, what does this indicate?

- A) Strong positive correlation
- B) Strong negative correlation
- C) No correlation
- D) Weak negative correlation